



Preparation & storage

Cow & Gate First Infant Milk, 800g EaZypack

Powder

| Approx. age | Approx. weight kg / lb | | No. of feeds per 24 hours | No. of level scoops per feed (1 scoop = 4.5g) | Single Feed ml / fl oz | |
|---------------|------------------------|--------|---------------------------|---|------------------------|-------|
| | kg | lb | | | ml | fl oz |
| Up to 2 weeks | 3.5 | 7 3/4 | 6 | 3 | 90 | 3 |
| 2-4 weeks | 3.9 | 8 1/2 | 5 | 4 | 120 | 4 |
| 4-8 weeks | 4.7 | 10 1/2 | 5 | 5 | 150 | 5 |
| 8-12 weeks | 5.4 | 12 | 5 | 6 | 180 | 6 |
| 3-4 months | 6.2 | 13 3/4 | 5 | 6 | 180 | 6 |
| 4-5 months | 6.9 | 15 1/4 | 5 | 7 | 210 | 7 |
| 5-6 months | 7.6 | 16 3/4 | 5 | 7 | 210 | 7 |
| 7-12 months | - | - | 3 | 7 | 210 | 7 |



1 Wash hands and sterilise all utensils according to manufacturers' instructions



2 Boil 1 litre of freshly run water. Leave to cool for 30 minutes. Do not use artificially softened or repeatedly boiled water.



3 Measure the required amount of water (refer to feeding guide) into a sterilised bottle. Do not add powder to boiling water due to risk of scalding.



4 Using the scoop provided, level off the powder with the built-in leveller. Do not press/heap the powder.



5 Add one scoop to every 30ml (1 fl oz) of boiled cooled water. Add the correct number of scoops to the water. Adding too many or too few scoops can be harmful.



6 Replace seal on bottle. Shake well (for 10 seconds) to dissolve powder. Remove seal, replace with a sterilised teat.



7 Cool under running tap. Check temperature of feed. Feed immediately.